

()

*

(/ / : / / :)

/ /

/ / /

°C

°C

/ /

%

/ meq/1000g / meq/1000g

/ mg/100g / mg/100g

× cfu/gr × cfu/gr

/ mg/100g

/ meq/1000g / meq/1000g

× cfu/gr

/ mg/100g

× cfu/gr

%

°C

:

.()

/ /

/ /

.()

.()

.()

/

/

FAO

.()

/

/

/

/

/

()

/

.()

.()

/

°C

°PV, °TVN ()

°BHA

/

/

()	(%)	(%)
/	BHA	B
/	BHA	B
/	BHA	B'
/	BHA	B'
/	CA	C
/	CA	C
/	CA	C'
/	CA	C'
		()S
/	BHA	B''
/	BHA	B''

Buchi 810
 Behr Labo-Technik GmbH
 COBOS-C-300-SX
 WF- Type-E-115 Heraeus
 Tuttlingen 7200 Binder

T.V.N
 .()
 .()
 .()

% / / BAH
 / /

3 . Total Volatile Nitrogen
 4 . Peroxid value

1. Clupeonella Ingrauliformis
 2 . Butylated hydroxyl anisol

meq/1000g
(°C)

/ f	/ e ^f	/ ef	/ e	/ d	/ C	/ b	a /	B
/ f	/ ef	/ e	/ e	/ cd	/ bc	/ ab	a /	B
/ e ^f	/ e	/ e	/ e	/ cd	/ b	a /	a /	B'
/ de	/ cd	/ cd	/ cd	/ c	/ b	a /	a /	B'
/ O	/ m	/ m	/ k	/ hi	/ d	/ c	a /	C
/ P	/ m	/ lm	/ j	/ h	/ cd	/ bc	a /	C
n	/ l	/ j	/ g	/ f	/ cd	/ b	a /	C'
/ n	/ hi	/ h	/ g	/ ef	/ c	/ b	a /	C'
/ hi	/ q	/ s	r	/ m	/ d	/ c	a /	()S
		/ l	/ g	/ e	/ d	/ b	/ ab	B''
		/ hi	/ fg	/ e	/ cd	/ ab	/ ab	B''

() ... , b, a

T.V.N

(°C)

/ l	/ h	/ fg	e	/ d	/ c	/ bc	/ a	B
i	/ gh	e	/ de	/ d	/ c	/ b	/ a	B
/ hi	/ g	/ fg	/ de	/ d	/ c	/ bc	/ a	B'
/ gh	/ fg	/ ef	/ de	/ cd	/ bc	/ b	/ a	B'
/ g	/ fg	/ e	/ de	/ d	/ bc	/ b	/ a	C
/ g	/ f	e	d	/ cd	/ bc	/ b	/ a	C
/ fg	/ f	e	/ d	/ cd	/ bc	/ b	/ a	C'
/ fg	/ ef	/ ef	/ de	/ cd	/ b	/ b	/ a	C'
/ j	/ i	/ h	/ fg	/ de	/ c	/ c	/ a	()S
		/ fg	e	/ cd	/ c	/ b	/ a	B''
		e	/ d	/ c	/ bc	/ b	/ a	B''

() ... , b, a

:

Cfu/gr

(°C)

x	a	x	a	x	a	x	ab	x	b	x	c	x	cd	B
x	a	x	a	x	a	x	b	x	bc	x	c	x	c	B
x	a	x	a	x	b	x	b	x	bc	x	bc	x	c	B'
x	a	x	a	x	a	x	b	x	bc	x	bc	x	c	B'
x	a	x	a	x	a	x	a	x	ab	x	bc	x	bc	C
x	a	x	a	x	ab	x	b	x	b	x	bc	x	bc	C
x	a	x	a	x	a	x	b	x	bc	x	bc	x	bc	C'
x	a	x	a	x	a	x	ab	x	b	x	b	x	bc	C'
x	b	x	e	x	f	x	f	x	g	x	g	x	h	()S
				x	a	x	ab	x	bc	x	c	x	d	B''
				x	a	x	a	x	bc	x	bc	x	cd	B''

()

..., b, a

B''	B''	C'	C'	C	C	B'	B'	B	B
/						/	/	/	/
/						/	/	/	/
/		/		/	/				
/	/		/						
/	/			/					

T.V.N
()
T.V.N

T.V.N , P.V

T.V.N

T.V.N

T.V.N

meq/ g /
meq/ g /

T.V.N

T.V.N

BHA

pH

pH

g /

T.V.N

meq/

/ C'
/

T.V.N

BHA

T.V.N

()

meq/ g / B'

()

Losada () Sreenivasan

() Turner ()

pH

Losada ()

()

B' B' B B

BHA

BHA
/ (GRAS)
/ BHA () ()
() (°C)

() Aitken pH BHA
Soniapedro ()
()

() ()
() ()

P.V T.V.N

()
" ()

/ BHA :B₁
/ BHA :B₂-₃
/ BHA :B'₁
3/ BHA :B'₂

5 . General Recognized As Safe

REFERENCES

7. Aitken, A. 2001. Polyphosphates in Fish Processing. Ministry of Agriculture, Fisheries and Food, TORRY Research Stations England. Advisory Note: No. 31:1-4.
8. Bracco, U. and et al., 1981. Production and use of Natural Antioxidants, J. A. Q. C. S. June.689-690.
9. Connell, J. J. 1997. Control of Fish Quality. Fishing News Books LTd. Farnham, Surrey, England, pp.37-38.
10. Dyer, W. J. 1969.Effect of brining and polyphosphate on yield and quality. Technology of Fish utilisation Ed. R. Kreuzer, Fishing News (book) LTd.,London, England pp. 167-179.
11. Dyer, W. J. 1971. Speed of Freezing and quality of Frozen Fish. Fish Inspection and Quality Control. Fishing News books, Farnham, p. 101.
12. FAO, 1978. Specifications for Identity and purity. Food and Agriculture Organization of the United Nations. Vol. 4: 15
13. Freeman, D. W., Shannon, C. W. 1994. Rancidity in Frozen Catfish Fillets as influenced by antioxidant injection and storage. J. - Aqual. - Food - Prodo. Technol. Vol. 3. no. 1: 65-76.
14. Foulke, J. E. 2003. A Fresh look at Preservatives. (internet), 2 pp.
15. Govindan, T. K. 1985. Fish Processing Technology. Oxford and IBH publishing Co. PVI.Lid NewDelhi, Bombay, Calcuita, pp. 89-91.
16. Hiremath. G. G. and Sreenivasan. N. 1979. Studies on prevention of quality loss in frozen seerfillets during storage by use of additives. J. Agric. Sci, 13: 88-92.
17. Hall, G. M. 1992. Fish Proccsing Technology. Publisher: Blackie Academic and Profesional. New York. London. Glaskow, pp. 31-32.
18. Harrigan, W. F. and Mc Cane, M. E. 1990. Laboratory Methods in Microbiology. Academic press, London and New York. pp. 52-53.
19. Lovern, J. A. 1962. The lipid of fish and changes occurring in them during processing and storage.In Fish in Nutrition, ed. E.Hean and R. Kreuzer. London: Fishing News LTd, pp. 35-37.

- :
20. Losada, L. 2003. Effect of citric acid on Rancidity Development in Horse mackerel during frozen storage. Department de Quimica de productos marinos, Eduardo cobello 6, E – 36208 vgo-Spain, (internet) , p. 1.
 21. Moini, S. 1980. Changes occurring in some chemical and sensory properties of smoked cod during cold storage. Ph.D. Thesis, Reading University, Reading, England, pp 220.
 22. Pedro. S. And et al., 2003. Cod spoilage Flora inhibition by citric acid and potassium sorbate. Brasilia, Portugal, (internet), p. 1.
 23. Sutton, A. H. , 1969. Polyphosphate Treatment of cod muscle. Technology of Fish utilization. Ed., R. Kreuzer, Fishing News (Books) LTd., London. pp. 172-176.
 24. Sreenivasan, N. and et al. 1976. Studies on Changes in Indian Mackerel during frozen storage and on the efficacy of protective Treatments in Inhibiting Rancidity. J. Agric. Sci. 10: 290-305.
 25. Turner. A. G. and et al. 2003. The Efficacy of using natural antioxidant blends to control Oxidative Rancidity in Minced Rainbow trout during frozen storage, Department of Food Science and Technology Duck pond Drive, Virginia Polytechnic Institute and State University.