

()

(Bombyx mori)

*

(/ / : / / :)

x

(P<%)

:

/

/

/

/

()

/

()

()

()

)

(

)

(
()

()

()

()

)

(

()

()

x

±

±

)

()

(

() () ()
()

x

()

()

%

... :

$$\frac{(C_{w2}-C_{w1})}{N_c}$$

()

P_b

W_b

()

T_f

|

()

$(F_{i2}-F_{i1})$

y

x

L_d

WS_h

SPSS

($P < \%$)

(C)

(B)

$$W_b T_f (F_{i2}-F_{i1}) L_d WS_h = B$$

$$(C_{w2}-C_{w1}) N_c P_b = C$$

()

% /

(± SE)

/ a	/ b	/ b	/ c	()
(± /)	(± /)	(± /)	(± /)	

/ a	/ b	/ bc	/ c	()
(± /)	(± /)	(± /)	(± /)	

/ a	/ a	/ a	/ a	
(± /)	(± /)	(± /)	(± /)	

/ a	/ a	/ a	/ a	
(± /)	(± /)	(± /)	(± /)	

(P<%)

/

/

/

/

/

/

/ a	/ b	/ c	/ d	()
-----	-----	-----	-----	-----

/ a	/ b	/ c	/ d	()
-----	-----	-----	-----	-----

/ a	/ ab	/ bc	/ d	(%)
-----	------	------	-----	-----

/ a	/ b	/ c	/ d	(%)
-----	-----	-----	-----	-----

/ c	/ c	/ b	/ a	
-----	-----	-----	-----	--

/ d	/ c	/ b	/ a	
-----	-----	-----	-----	--

/ c	/ bc	/ b	/ a	
-----	------	-----	-----	--

/ d	/ c	/ b	/ a	
-----	-----	-----	-----	--

(P<%)

() / /

/

% /

% /

/

...

:

()

(+)

()

(/)

(/)

(/)

+

+

+

/

/

/

/

/

/

()

,

()

()

()

+

)

/ (

/ ()

.() %

.()

.() ()

.() ()

/ / .() /
/ / / /

()

.()

... :
%
%
()
% / % /
% /
()

REFERENCES

(*Bombyx mori* L.)
Bombyx mori L.

6. Bora B., T.S., Mahadevamurthy, G.R., Shivakumar, & S.B. Magadam. 1995. Effect of feed quantum during fifth age silkworm, *Bombyx mori* L. on reeling parameters. *Uttar Pradesh J. Zoo.* 15(1): 1-6.
7. Broadway, R.M. & S.S., Duffey. 1986. The effect of dietary protein on the growth and digestive physiology of larval *Heliothis zea* and *Spodoptera exigua*. *J. Insect Physiol.*, 32(8): 673-680.
8. Chapman, R.F. 1998. *The Insect Structure and Function*, 4th edition, Cambridge University Press, Cambridge.
9. Das, B.K., J. Sarker, C. Das, N. K. Das, & S.K., Sen. 1995. Seasonal effects on the relative performance of five biovoltine breeds of silkworm *Bombyx mori*. *Uttar Pradesh J. Zoo.* 15(2): 91-96.

10. Das S., M. Shamsuddin, & K. Sengupta. 1991. Minimization of feeding frequency has no adverse effect of the bivoltine crop production in West Bengal. *Indian Silk*, 30(8):30.
11. Das S., P. K. Saha, M. Shamsudin, & S.K. Sen. 1994. Feeding frequency: The economic potentiality and efficacy in tropical bivoltine rearing during the favorable season. *Sericologia*, 34(3):533-536.
12. Etebari, K., R. Ebadi, & L. Matindoost. 2004. Effect of feeding mulberry's enriched leaves with ascorbic acid on some biological, biochemical and economical characteristics of silkworm *Bombyx mori* L. *Int. J. Indust. Entomol.* 8(1): 81-87.
13. Fonseca, T.C., J. E. Almedia, & A.S. Fonseca. 1990. Effect of mulberry selection on silkworm feeding. *Sericologia*, 30(4): 475-477.
14. Giridhar, K. & N.S., Reddy. 1991. Effective rate of rearing in bivoltine silkworm, *Bombyx mori* breeds on different mulberry, *Morus* sp. varieties. *Indian J. Seric.*, 30(1):88-90.
15. He G., Wu P. & Y. Huang. 1987. Economic efficiency of scarce rearing of *Bombyx mori*. *Guangdong Agric Sci.* 3:23-25.
16. Hof, H.M., & M. Martin. 1989. The effect of diet water content on energy expenditure by third instar *Manduca sexta* larvae (Lepidoptera: Sphingidae). *J. Insect Physiol.*,35(5): 433-436.
17. Ito, T. 1978. Silkworm Nutrition, In "*The Silkworm an Important Laboratory Tools*", Tazima, Y., (ed), Kodansha Ltd., Tokyo, pp. 121-157.
18. Koilpillai, R.R. 1995. Impact of nutritional modulation on survival, growth, bioenergetics and reproductive potential in the silkworm, *Bombyx mori* L. 19(3) 223-227.
19. Koilpillai, R.R. 1995. Morphometric studies on the silkworm, *Bombyx mori* (L.) as a function of feeding duration. 19(4): 321-328.
20. Krishnaswami S., M. N. Narasimhanna, S. K. Suryanarayan, & S. Kumararaj. 1973. Sericulture manual,2, silkworm rearing,FAO Agricultural Services Bulletine.
21. Li, R., & Q. Sang. 1984. The relationship between quality of mulberry leaves and some economic characters during the later larval stage. *Science of Sericulture Canye Kexue.* 10(4):197-201. (In Chienes with English summery)
22. Lim, S.H., Y. T. Kim, S. P. Lee, I. J. Rhee, J. S. Lim, & B.H., Lim. 1990. Sericulture training manual, FAO, Agricultural Services Bulletin, Rome, p. 103.
23. Lu, X.F. 1992. Relation between mulberry feeding amounts and cocoon weight, cocoon yield and filament quality in three moltine cocoon production, *Acta Sericologica Sinica*, 18(1): 49-52. (In Chienes with English summery).
24. Maheshkumar Vage N. & J. Ashoka. 2000. Effect of tender shoot feeding on silk technological parameters of silkworm, *Bombyx mori*. *Sericologia* 40(1): 79-89.
25. Meenal A. & O. Ninagi. 1995. Comparative study of ingestion pattern in polyvoltine and bivoltine silkworm races at different feeding levels. *Sericologia*, 35(4): 747-752.
26. Mukerji, M.K. & J.C. Guppy. 1970. A quantitative study of food consumption and growth in *Pseudaletia unipuncta* (Lep., Noctuidae), *Can. Ent.*, 102,1179-1188.
27. Narayanan E.S. & S. S. Chawla. 1965. Effect of frequency of feeding on the growth and development in two races of silkworm. *Indian J. Seri.*4(4): 1-3.
28. Paul, D.C, G. S. Rao, & D.C. Deb. 1992. Impact of dietary moisture on nutritional indices and growth of *Bombyx mori* and concomitant larval duration, *J. Insect Physiol.*, 38(3): 229-235.
29. Raman K.V., S.B. Magadum & R.K. Datta. 1994. Feed efficiency of the silkworm *Bombyx mori* L. hybrid (NB4DD X KA). *Insect Sc. Appl.* 15(2): 111-116.
30. Remadevi O.K., S. B. Magadum, K. V. Benchamin & R.K. Datta. 1993. Mutual correlation among the nutritional and economic characters of the multivoltine silkworm, *Bombyx mori* L. *Indian J. Seri.*32(2): 189-195.

31. Singh G.B. & O. Ninagi. 1995. Comparative studies on food utilization efficiency in some silkworm strains under different feeding levels. *Sericologia*, 35(4): 667-675.
32. Sharma, D.K. & D. Devi. 1997. Seasonal variation of the foliar constituents of the primary food plants of the muga silkworm (*Antheraea assama* WW.). *Sericologia*, 37(2): 251-258.
33. Verma, S.K. & K.S., Kushwaha. 1970. Comparative growth of the silkworm, *Bombyx mori* race Bulupolu reared on different mulberry varieties. *Indian J. Agric. Sci.*, 40(12): 1097-1107.
34. Volney, W.J., J. E. Milstead, & V. R. Lewis. 1983. Effect of food quality, larval density, and photoperiod on the feeding rate of the california oakworm (Lep., Diptidae). *Environ. Entomol.*, 12, 792-798.
35. Waldbauer, G.P. 1968. the consumption and utilization of food by insects. *Adv. Insect Physiol.* 5: 229-288.
36. Woods, H.A., & J.G. Kingsolver. 1999. Feeding rate and the structure of protein digestion and absorption in lepidopteran midgut *Arch. Insect Biochem. Physiol.*, 42: 74-87.